FORMAL SELF-ADJOINTNESS OF A FAMILY OF CONFORMALLY INVARIANT BIDIFFERENTIAL OPERATORS

JEFFREY S. CASE AND ZETIAN YAN

ABSTRACT. We prove that the curved Ovsienko–Redou operators and a related family of differential operators are formally self-adjoint. This verifies two conjectures of Case, Lin, and Yuan.

1. Introduction

The GJMS operator of order 2k, introduced by Graham, Jenne, Mason, and Sparling [7], is a conformally invariant differential operator with leading-order term Δ^k defined on any Riemannian manifold (M^n, g) of dimension $n \geq 2k$. It can be defined by suitably restricting the power $\tilde{\Delta}^k$ of the Laplacian of the Fefferman–Graham ambient space [4]. While $\tilde{\Delta}^k$ is formally self-adjoint in the ambient space, this does not immediately imply that the GJMS operators are formally self-adjoint.

There are now two proofs that the GJMS operators are formally self-adjoint. The first, due to Graham and Zworski [8] (cf. [3]), uses a realization of the GJMS operators as poles of the scattering operator of a Poincaré space. The second, due to Juhl [9] (cf. [5]), uses a remarkable formula for the GJMS operators in terms of a family of formally self-adjoint second-order differential operators. The first approach relies heavily on the equivalent characterization of the GJMS operators as obstructions to formally extending a function to be harmonic in the ambient space, while the second approach requires a complicated combinatorial argument.

Case, Lin, and Yuan [2] gave two generalizations of the GJMS operators; here we discuss those which are plausibly formally self-adjoint. First is a family of conformally invariant bidifferential operators

$$D_{2k} \colon \mathcal{E}\left[-\frac{n-2k}{3}\right]^{\otimes 2} \to \mathcal{E}\left[-\frac{2n+2k}{3}\right]$$

of total order 2k. They are called the **curved Ovsienko–Redou operators** because they generalize a family of bidifferential operators constructed by Ovsienko and Redou [10] on the sphere. The operators D_{2k} are determined ambiently by

$$\begin{split} \widetilde{D}_{2k}(\widetilde{u}\otimes\widetilde{v}) := \sum_{r+s+t=k} a_{r,s,t} \widetilde{\Delta}^r \left((\widetilde{\Delta}^s \widetilde{u}) (\widetilde{\Delta}^t \widetilde{v}) \right), \\ a_{r,s,t} := \frac{k!}{r!s!t!} \frac{\Gamma\big(\frac{n+4k}{6} - r\big) \Gamma\big(\frac{n+4k}{6} - s\big) \Gamma\big(\frac{n+4k}{6} - t\big)}{\Gamma\big(\frac{n-2k}{6}\big) \Gamma\big(\frac{n+4k}{6}\big)^2}, \end{split}$$

 $^{2020\} Mathematics\ Subject\ Classification.\ Primary\ 53C25.$

Key words and phrases. conformally invariant operators; Ovsienko–Redou operators; formal self-adjointness.

on $\widetilde{\mathcal{E}}\left[-\frac{n-2k}{3}\right]\otimes\widetilde{\mathcal{E}}\left[-\frac{n-2k}{3}\right]$; in this paper, tensor products are over \mathbb{R} . Second is a family of conformally invariant differential operators

$$D_{2k,\mathcal{I}} \colon \mathcal{E}\left[-\frac{n-2k-2\ell}{2}\right] \to \mathcal{E}\left[-\frac{n+2k+2\ell}{2}\right]$$

of order 2k associated to a scalar Weyl invariant \mathcal{I} of weight -2ℓ . These are determined ambiently by a scalar Riemannian invariant \widetilde{I} of weight -2ℓ and

$$\begin{split} \widetilde{D}_{2k,\widetilde{I}}(\widetilde{u}) &:= \sum_{r+s=k} b_{r,s} \widetilde{\Delta}^r \left(\widetilde{I} \widetilde{\Delta}^s \widetilde{u} \right), \\ b_{r,s} &:= \frac{k!}{r!s!} \frac{(\ell+s-1)!(\ell+r-1)!}{(\ell-1)!^2}, \end{split}$$

on $\widetilde{\mathcal{E}}\left[-\frac{n-2k-2\ell}{2}\right]$. See Section 2 for an explanation of our notation and a description of how the ambient formulas determine conformally invariant operators.

The symmetry of the coefficients $a_{r,s,t}$ and $b_{r,s}$ and the observations

$$\widetilde{\mathcal{E}}\left[-\frac{n-2k}{3}\right] \ni \widetilde{u}, \widetilde{v}, \widetilde{w} \Longrightarrow \widetilde{u}\widetilde{D}_{2k}(\widetilde{v} \otimes \widetilde{w}) \in \widetilde{\mathcal{E}}[-n],$$

$$\widetilde{\mathcal{E}}\left[-\frac{n-2k-2\ell}{2}\right] \ni \widetilde{u}, \widetilde{v} \Longrightarrow \widetilde{u}\widetilde{D}_{2k,\widetilde{I}}(\widetilde{v}) \in \widetilde{\mathcal{E}}[-n],$$

imply that the induced operators D_{2k} and $D_{2k,\mathcal{I}}$ are plausibly formally self-adjoint. Case, Lin, and Yuan [2] conjectured their formally self-adjointness and verified this when $k \leq 3$. The difficulty in this problem is that there is not an equivalent description of these operators as an obstruction to solving some second-order PDE, and hence the Graham–Zworski argument cannot be adapted to D_{2k} or $D_{2k,\mathcal{I}}$. Case, Lin, and Yuan instead adapted Juhl's approach, with the constraint $k \leq 3$ due to the difficulty of the combinatorial argument.

In this paper we develop a new, conceptually simple approach to proving formal self-adjointness which verifies the Case–Lin–Yuan conjecture:

Theorem 1.1. Let (M^n, g) be a Riemannian manifold and let $k \in \mathbb{N}$; if n is even, then assume additionally that $k \leq n/2$. Then $D_{2k} \colon \mathcal{E}\left[-\frac{n-2k}{3}\right]^{\otimes 2} \to \mathcal{E}\left[-\frac{2n+2k}{3}\right]$ is formally self-adjoint.

Theorem 1.2. Let (M^n, g) be a Riemannian manifold, let $k \in \mathbb{N}$, and let \mathcal{I} be a scalar Weyl invariant of weight -2ℓ in the ambient space; if n is even, then assume additionally that $k \leq n/2$ if $\ell = 0$ and $k + \ell \leq n/2 + 1$ if $\ell \geq 1$. Then $D_{2k,\mathcal{I}} \colon \mathcal{E}\left[-\frac{n-2k-2\ell}{2}\right] \to \mathcal{E}\left[-\frac{n+2k+2\ell}{2}\right]$ is formally self-adjoint.

The restrictions on k and $k + \ell$ when n is even ensure that the operators D_{2k} and $D_{2k,\mathcal{I}}$, respectively, are independent of the ambiguity of the ambient metric; see Section 2 for details. When $\mathcal{I} = 1$, the operator $D_{2k,\mathcal{I}}$ is a nonzero multiple of the GJMS operator of order 2k, and so Theorem 1.2 gives a new proof of its formal self-adjointness.

The main idea in the proofs of Theorems 1.1 and 1.2 is to restrict the ambient operators \widetilde{D}_{2k} and $\widetilde{D}_{2k,\widetilde{I}}$ to the Poincaré space (\mathring{X},g_+) equivalent to a given ambient space. Their Dirichlet forms

$$\int_{r>\varepsilon} \left. \left(\widetilde{u} \widetilde{D}_{2k} (\widetilde{v} \otimes \widetilde{w}) \right) \right|_{\mathring{X}} \mathrm{dV}_{g_+} \quad \text{ and } \quad \int_{r>\varepsilon} \left. \left(\widetilde{u} \widetilde{D}_{2k,\widetilde{I}} (\widetilde{v}) \right) \right|_{\mathring{X}} \mathrm{dV}_{g_+}$$

are both in $O(\log \varepsilon)$, and the coefficient of the log terms are the Dirichlet forms of the respective operators D_{2k} and $D_{2k,\mathcal{I}}$ on M. The Divergence Theorem implies that there are polydifferential operators B and $B_{\widetilde{I}}$ such that

$$\int_{r>\varepsilon} \left(\widetilde{u} D_{2k}(\widetilde{v} \otimes \widetilde{w}) - \widetilde{v} \widetilde{D}_{2k}(\widetilde{u} \otimes \widetilde{w}) \right) \Big|_{\mathring{X}} dV_{g_{+}} = \int_{r=\varepsilon} B(u \otimes v \otimes w) dA_{g_{+}},$$

$$\int_{r>\varepsilon} \left(\widetilde{u} \widetilde{D}_{2k,\widetilde{I}}(\widetilde{v}) - \widetilde{v} \widetilde{D}_{2k,\widetilde{I}}(\widetilde{u}) \right) \Big|_{\mathring{X}} dV_{g_{+}} = \int_{r=\varepsilon} B_{\widetilde{I}}(u \otimes v) dA_{g_{+}}.$$

Moreover, the right-hand sides cannot have a log term—they are in O(1)—and hence D_{2k} and $D_{2k,\mathcal{I}}$ are formally self-adjoint; see Section 4 for details.

This paper is organized as follows:

In Section 2 we recall necessary background material on the ambient space and give a unified construction of the operators D_{2k} and $D_{2k,\mathcal{I}}$ of Theorem 1.1 and Theorem 1.2, respectively. Our construction differs slightly from that of Case, Lin, and Yuan. We normalize our operators so that D_{2k} coincides with their construction of the curved Ovsienko–Redou operators. While the operators $D_{2k,\mathcal{I}}$ do not coincide with the linear operators constructed by Case, Lin, and Yuan [2, Theorem 1.7], the linear spans of our respective operators agree.

In Section 3 we describe the restrictions of \widetilde{D}_{2k} and $\widetilde{D}_{2k,\widetilde{I}}$ to (\mathring{X},g_+) and express them in terms of linear combinations of compositions of $\Delta_{g_+}+c$ for constants $c\in\mathbb{R}$.

In Section 4 we realize the Dirichlet forms for D_{2k} and $D_{2k,\mathcal{I}}$ in terms of the leading-order term of the corresponding Dirichlet forms in (\mathring{X}, g_+) . We then use the simple formulas for the latter operators to deduce Theorems 1.1 and 1.2.

2. Ambient spaces and the Ovsienko-Redou operators

In this section we recall the construction [2] of the curved Ovsienko–Redou operators and their related linear operators.

2.1. **Ambient spaces.** We begin by recalling the relevant aspects of the ambient space, following Fefferman and Graham [4].

Let (M^n, \mathfrak{c}) be a conformal manifold of signature (p, q). Denote

$$\mathcal{G} := \{(x, q_x) : x \in M, q \in \mathfrak{c}\} \subset S^2 T^* M$$

and let $\pi: \mathcal{G} \to M$ be the natural projection. We regard \mathcal{G} as a principal \mathbb{R}_+ -bundle with dilation $\delta_{\lambda}: \mathcal{G} \to \mathcal{G}, \ \lambda \in \mathbb{R}_+$, given by

$$\delta_{\lambda}(x, g_x) := (x, \lambda^2 g_x).$$

Denote by $T := \frac{\partial}{\partial \lambda}|_{\lambda=1} \delta_{\lambda}$ the infinitesimal generator of δ_{λ} . The canonical metric is the degenerate metric g on \mathcal{G} defined by

$$\boldsymbol{g}(X,Y) := g_x(\pi_* X, \pi_* Y)$$

for $X, Y \in T_{(x,q_x)}\mathcal{G}$. Note that $\delta_{\lambda}^* \mathbf{g} = \lambda^2 \mathbf{g}$.

A choice of representative $g \in \mathfrak{c}$ determines an identification $\mathbb{R}_+ \times M \cong \mathcal{G}$ via $(t,x) \cong (x,t^2g_x)$. In these coordinates, $T_{(t,x)} = t\partial_t$ and $g_{(t,x)} = t^2\pi^*g$.

Extend the projection and dilation to $\mathcal{G} \times \mathbb{R}$ in the natural way:

$$\pi(x, g_x, \rho) := x,$$

$$\delta_{\lambda}(x, g_x, \rho) := (x, \lambda^2 g_x, \rho),$$

where ρ denotes the coordinate on \mathbb{R} . We abuse notation and also denote by T the infinitesimal generator of $\delta_{\lambda} \colon \mathcal{G} \times \mathbb{R} \to \mathcal{G} \times \mathbb{R}$. Let $\iota \colon \mathcal{G} \to \mathcal{G} \times \mathbb{R}$ denote the inclusion $\iota(x,g_x) := (x,g_x,0)$. A **pre-ambient space** for (M^n,\mathfrak{c}) is a pair $(\widetilde{\mathcal{G}},\widetilde{g})$ consisting of a dilation-invariant subspace $\widetilde{\mathcal{G}} \subseteq \mathcal{G} \times \mathbb{R}$ containing $\iota(\mathcal{G})$ and a pseudo-Riemannian metric \widetilde{g} of signature (p+1,q+1) satisfying $\delta_{\lambda}^*\widetilde{g} = \lambda^2\widetilde{g}$ and $\iota^*\widetilde{g} = \mathbf{g}$.

An **ambient space** for (M^n, \mathfrak{c}) is a pre-ambient space $(\widetilde{\mathcal{G}}, \widetilde{g})$ for (M^n, \mathfrak{c}) which is formally Ricci flat; i.e.

- if n is odd, then $\operatorname{Ric}(\widetilde{g}) \in O(\rho^{\infty})$;
- if n is even, then $\operatorname{Ric}(\widetilde{g}) \in O^+(\rho^{n/2-1})$.

Here $O^+(\rho^m)$ is the set of sections S of $S^2T^*\widetilde{\mathcal{G}}$ such that

- (i) $\rho^{-m}S$ extends continuously to $\iota(\mathcal{G})$, and
- (ii) for each $z = (x, g_x) \in \mathcal{G}$, there is an $s \in S^2T_x^*M$ such that $\operatorname{tr}_{g_x} s = 0$ and $(\iota^*(\rho^{-m}S)(z) = (\pi^*s)(z)$.

Two ambient spaces $(\widetilde{\mathcal{G}}_j, \widetilde{g}_j)$, $j \in \{1, 2\}$, for (M^n, \mathfrak{c}) are **ambient-equivalent** if there are open sets $\mathcal{U}_j \subseteq \widetilde{\mathcal{G}}_j$ and a diffeomorphism $\Phi \colon \mathcal{U}_1 \to \mathcal{U}_2$ such that

- (i) $\iota(\mathcal{G}) \subseteq \mathcal{U}_j$ for each $j \in \{1, 2\}$;
- (ii) each \mathcal{U}_j is dilation-invariant and Φ commutes with dilations;
- (iii) $\Phi \circ \iota = \iota$;
- (iv) if n is odd, then $\Phi^*\widetilde{g}_2 \widetilde{g}_1 \in O(\rho^{\infty})$;
- (v) if n is even, then $\Phi^* \widetilde{g}_2 \widetilde{g}_1 \in O^+(\rho^{n/2})$.

Fefferman and Graham showed [4, Theorem 2.3] is that if (M^n, \mathfrak{c}) is a conformal manifold, then there is a unique, up to ambient-equivalence, ambient space for (M^n, \mathfrak{c}) . In fact, they proved a stronger statement [4, Theorem 2.9(A)]: Let (M^n, \mathfrak{c}) be a conformal manifold and pick a representative $g \in \mathfrak{c}$. Then there is an $\varepsilon > 0$ and a one-parameter family g_ρ , $\rho \in (-\varepsilon, \varepsilon)$, of metrics on M such that $g_0 = g$ and

(2.1)
$$\widetilde{\mathcal{G}} := \mathcal{G} \times (-\varepsilon, \varepsilon),$$

$$\widetilde{g} := 2\rho dt^2 + 2t dt d\rho + t^2 g_\rho$$

defines an ambient space $(\mathcal{G}, \mathcal{g})$ for (M^n, \mathfrak{c}) . We say that an ambient metric in the form above is **straight and normal**.

Let $(\widetilde{\mathcal{G}}, \widetilde{g})$ be the ambient space for (M^n, \mathfrak{c}) . Denote by

$$\widetilde{\mathcal{E}}[w] := \left\{ \widetilde{f} \in C^{\infty}(\widetilde{\mathcal{G}}) : \delta_{\lambda}^{*} \widetilde{f} = \lambda^{w} \widetilde{f} \right\}$$

the space of homogeneous functions on $\widetilde{\mathcal{G}}$ of weight $w \in \mathbb{R}$. Note that $\widetilde{f} \in \widetilde{\mathcal{E}}[w]$ if and only if $T\widetilde{f} = w\widetilde{f}$. The space of **conformal densities** of weight w is

$$\mathcal{E}[w] := \left\{ \iota^* \widetilde{f} \in C^\infty(\mathcal{G}) : \widetilde{f} \in \widetilde{\mathcal{E}}[w] \right\}.$$

Fix $n \in \mathbb{N}$. An ambient scalar Riemannian invariant \widetilde{I} is an assignment to each ambient space $(\widetilde{\mathcal{G}}^{n+2}, \widetilde{g})$ of a linear combination $\widetilde{I}_{\widetilde{g}}$ of complete contractions of

(2.2)
$$\widetilde{\nabla}^{N_1} \widetilde{\operatorname{Rm}} \otimes \cdots \otimes \widetilde{\nabla}^{N_\ell} \widetilde{\operatorname{Rm}},$$

with $\ell \geq 2$, where $\widetilde{\nabla}$ and $\widetilde{\operatorname{Rm}}$ are the Levi-Civita connection and Riemann curvature tensor, respectively, of \widetilde{g} , we regard $\widetilde{\operatorname{Rm}}$ as a section of $\otimes^4 T^* \widetilde{\mathcal{G}}$, and we use \widetilde{g}^{-1} to

take contractions. Any complete contraction of (2.2) is homogeneous of weight

$$w = -2\ell - \sum_{i=1}^{\ell} N_i.$$

We assume $\ell \geq 2$ because any complete contraction of $\widetilde{\nabla}^N \widetilde{\mathrm{Rm}}$ is proportional to $\widetilde{\Delta}^{N/2}\widetilde{R}$ modulo ambient scalar Riemannian invariants, and $\widetilde{\Delta}^{N/2}\widetilde{R}=0$ when it is independent of the ambiguity of \widetilde{g} . If \widetilde{I} is independent of the ambiguity of \widetilde{g} , then $\mathcal{I} := \iota^* \widetilde{I}_{\widetilde{a}} \in \mathcal{E}[w]$ is independent of the choice of ambient space. A scalar Weyl **invariant** is a scalar invariant $\mathcal{I} \in \mathcal{E}[w]$ constructed in this way. Fefferman and Graham gave a condition on the weight w which implies this independence:

Lemma 2.1 ([4, Proposition 9.1]). Let $(\widetilde{\mathcal{G}}^{n+2}, \widetilde{g})$ be a straight and normal ambient space and let $\widetilde{I} \in \widetilde{\mathcal{E}}[w]$ be an ambient scalar Riemannian invariant. If $w \geq -n-2$, then $\iota^*\widetilde{I}_{\widetilde{g}}$ is independent of the ambiguity of \widetilde{g} .

Bailey, Eastwood, and Graham [1, Theorem A] showed that every conformally invariant scalar of weight w > -n is a Weyl invariant.

2.2. Conformally invariant polydifferential operators. Fix $k, n \in \mathbb{N}$. An ambient polydifferential operator D of weight -2k is an assignment to each ambient space $(\widetilde{\mathcal{G}}^{n+2}, \widetilde{g})$ of a linear map

$$\widetilde{D}^{\widetilde{g}} : \widetilde{\mathcal{E}}[w_1] \otimes \cdots \otimes \widetilde{\mathcal{E}}[w_j] \to \widetilde{\mathcal{E}}[w_1 + \cdots + w_j - 2k]$$

such that $\widetilde{D}^{\widetilde{g}}(\widetilde{u}_1 \otimes \cdots \otimes \widetilde{u}_i)$ is a linear combination of complete contractions of

$$(2.3) \qquad \widetilde{\nabla}^{N_1} \widetilde{u}_1 \otimes \cdots \otimes \widetilde{\nabla}^{N_j} \widetilde{u}_j \otimes \widetilde{\nabla}^{N_{j+1}} \widetilde{\operatorname{Rm}} \otimes \cdots \otimes \widetilde{\nabla}^{N_\ell} \widetilde{\operatorname{Rm}}$$

with $\ell = j$ or $\ell \geq j + 2$. Necessarily the powers N_1, \ldots, N_ℓ satisfy

$$\sum_{i=1}^{\ell} N_i + 2\ell - 2j = 2k.$$

The **total order** of such a contraction is $\sum_{i=1}^{j} N_i$. We say that \widetilde{D} is **tangential** if $\iota^*(\widetilde{D}^{\widetilde{g}}(\widetilde{u}_1 \otimes \cdots \otimes \widetilde{u}_j))$ depends only on $\iota^*\widetilde{u}_1, \ldots, \iota^*\widetilde{u}_j$ and \widetilde{g} modulo its ambiguity. On each conformal manifold (M^n, \mathfrak{c}) , such an operator determines a **conformally** invariant polydifferential operator

$$D: \mathcal{E}[w_1] \otimes \cdots \otimes \mathcal{E}[w_i] \to \mathcal{E}[w_1 + \cdots + w_i - 2k].$$

We will give a condition on the total order of an ambient polydifferential operator which implies that it is independent of the ambiguity of \widetilde{g} . Let $(\mathcal{G}, \widetilde{g})$ be a straight and normal ambient space for (M^n, g) . Following Fefferman and Graham [4], given coordinates $\{x^i\}_{i=1}^n$ on M and a multi-index $A=(a_1,\ldots,a_r)\in\{0,1,\ldots,n,\infty\}^r$, we denote by $\widetilde{T}_A = \widetilde{T}_{a_1 \cdots a_r}$ a component of a tensor $\widetilde{T} \in \otimes^k T^* \widetilde{\mathcal{G}}$, where the index 0 represents ∂_t , an index $i \in \{1, \ldots, n\}$ represents ∂_{x^i} , and the index ∞ represents ∂_{ρ} . The **length** of A is |A| := r and the **strength** of A is

$$||A|| := |\{j : a_j \in \{1, \dots, n\}\}| + 2 |\{j : a_j = \infty\}|.$$

When $\widetilde{f} \in C^{\infty}(\widetilde{\mathcal{G}})$, we denote $\widetilde{f}_{a_1 \cdots a_r} := \widetilde{\nabla}_{a_r} \cdots \widetilde{\nabla}_{a_1} \widetilde{f} =: (\widetilde{\nabla}^r \widetilde{f})_{a_r \cdots a_1}$. We first determine the dependence of the covariant derivatives $\widetilde{\nabla}^r \widetilde{f}$ of a function $\widetilde{f} \in C^{\infty}(\widetilde{\mathcal{G}})$ on the ambiguity of \widetilde{g} (cf. [4, Proof of Proposition 6.2]).

Lemma 2.2. Let $(\widetilde{\mathcal{G}}^{n+2}, \widetilde{g})$ be a straight and normal ambient space and let $\widetilde{f} \in \widetilde{\mathcal{E}}[w]$. Let A be a multi-index of length r := |A| and strength s := ||A||. Then \widetilde{f}_A mod $O(\rho^{(n-s)/2})$ depends only on \widetilde{g} mod $O(\rho^{n/2})$.

Proof. The proof is by induction in r. Since $\widetilde{f} \in \widetilde{\mathcal{E}}[w]$, we see that $\mathcal{L}_T \widetilde{\nabla}^r \widetilde{f} = w \widetilde{\nabla}^r \widetilde{f}$. Combining this with the identity $\widetilde{\nabla} T = \widetilde{g}$ yields

$$(2.4) \widetilde{\nabla}_T \widetilde{\nabla}^r \widetilde{f} = (w - r) \widetilde{\nabla}^r \widetilde{f}.$$

In particular, it suffices to consider the case $(n-s)/2 \le n/2-1$: If A is a multiindex of strength $s \le 1$, then Equation (2.4) implies that \widetilde{f}_A is proportional to \widetilde{f} (if s=0) or \widetilde{f}_a for some $a \in \{1,\ldots,n\}$ (if s=1). In either case \widetilde{f}_A is independent of \widetilde{g} .

We now proceed with the induction. The claim is trivially true when r = 0. For the inductive step, consider \widetilde{f}_{Aa} for $A = (a_1, \ldots, a_r)$ a multi-index of length r and $Aa := (a_1, \ldots, a_r, a)$ the corresponding multi-index of length r + 1. Denote s := ||Aa|| and s' := ||A||.

Suppose first that a = 0. Then Equation (2.4) yields $\widetilde{f}_{Aa} = (w - r)\widetilde{f}_A$. The required dependence of \widetilde{f}_{Aa} then follows from the inductive hypothesis.

Suppose next that $a \neq 0$. Write

$$\widetilde{f}_{Aa} = \partial_a \widetilde{f}_A - \widetilde{\Gamma}_{aa_1}^b \widetilde{f}_{ba_2 \cdots a_r} - \cdots - \widetilde{\Gamma}_{aa_r}^b \widetilde{f}_{a_1 \cdots a_{r-1}b}.$$

If $a \neq \infty$, then s = s' + 1 and hence $\partial_{x^i}O(\rho^{(n-s')/2}) \subseteq O(\rho^{(n-s)/2})$. If instead $a = \infty$, then s = s' + 2 and hence $\partial_{\rho}O(\rho^{(n-s')/2}) = O(\rho^{(n-s)/2})$. In either case, the inductive hypothesis then gives the required dependence of $\partial_a \widetilde{f}_A$. We now consider the summand $\widetilde{\Gamma}^b_{aa_r}\widetilde{f}_{a_1\cdots a_{r-1}b}$; the remaining summands are similar. Since $(n-s)/2 \leq n/2-1$, the formula [4, Equation (3.16)] for the Christoffel symbols of \widetilde{g} imply that $\widetilde{\Gamma}^K_{IJ}$ mod $O(\rho^{(n-s)/2})$ depends only on \widetilde{g} mod $O(\rho^{n/2})$, and so we can ignore the ambiguity caused by the Christoffel symbols. The worst case for the ambiguity caused by the inductive hypothesis is when $b = \infty$ and $a_r = 0$, so that $\|(a_1,\ldots,a_{r-1},b)\| = s' + 2$. If $a = \infty$, then s' + 2 = s, and so the required dependence follows from the inductive hypothesis. If $a \neq \infty$, then the vanishing of the Christoffel symbol Γ^∞_{a0} implies that the worst case does not happen. Thus $\|(a_1,\ldots,a_{r-1},b)\| \leq s' + 1 = s$, and so again the required dependence follows from the inductive hypothesis.

We now give a condition on the total order of an ambient polydifferential operator that implies that it is independent of the ambiguity of \tilde{g} (cf. [4, Proposition 9.1]).

Corollary 2.3. Let $(\widetilde{\mathcal{G}}^{n+2}, \widetilde{g})$ be a straight and normal ambient space and let \widetilde{D} be an ambient polydifferential operator of weight -2k. Suppose that

- (i) n is odd,
- (ii) $k \leq n/2$, or
- (iii) $k \leq n/2 + 1$ and D can be expressed as a linear combination of complete contractions of tensors of the form (2.3) with $\ell \geq j + 2$.

Then D is independent of the ambiguity of \tilde{g} .

Proof. The conclusion is clear when n is odd. Assume that n is even.

Since the component \tilde{g}^{ab} at $\rho = 0$ of the inverse ambient metric is nonzero only if ||ab|| = 2, we see that a complete contraction of a tensor of the form (2.3) can only

be nonzero on \mathcal{G} if each contracted pair has strength two. Denote by S_1, \ldots, S_ℓ the strengths of the factors in a contributing monomial. Note that the sum of the strengths must equal the total number of indices in a given monomial. Therefore

(2.5)
$$\sum_{i=1}^{\ell} S_i = \sum_{i=1}^{\ell} N_i + 4(\ell - j) = 2(k + \ell - j).$$

Clearly $S_i \geq 0$ if $i \leq j$. Fefferman and Graham observed [4, Proposition 6.1] that $S_i \geq 4$ if $i \geq j+1$.

Suppose first that $k \leq n/2$. If $i_0 \in \{1, ..., j\}$, then Equation (2.5) yields

$$S_{i_0} \le \sum_{i=1}^{\ell} S_i - 4(\ell - j) = 2(k - \ell + j) \le 2k \le n.$$

Moreover, equality holds if and only if k=n/2 and $\ell=j$ and $S_i=0$ for each $i\neq i_0$. If equality holds, then our monomial is proportional to $(\prod_{i\neq i_0}\widetilde{u}_i)\widetilde{\Delta}^{n/2}\widetilde{u}_{i_0}$ modulo complete contractions of tensors of the form (2.3) with $\ell\geq j+1$; these will be considered next. Graham et al. observed [7, Section 3] that $(\prod_{i\neq i_0}\widetilde{u}_i)\widetilde{\Delta}^{n/2}\widetilde{u}_{i_0}$ is independent of the ambiguity of \widetilde{g} . Otherwise Lemma 2.2 implies that the contribution of the factor $\widetilde{\nabla}^{M_{i_0}}\widetilde{u}_{i_0}$ is independent of the ambiguity of \widetilde{g} . If instead $i_0\in\{j+1,\ldots,\ell\}$, then $\ell\geq j+1$ and hence Equation (2.5) yields

$$S_{i_0} \le \sum_{i=1}^{\ell} S_i - 4(\ell - j - 1) = 2(k + 2 - \ell + j) \le n + 2.$$

Moreover, equality holds if and only if k = n/2 and $\ell = j+1$ and $S_i = 0$ for $i \neq i_0$. If equality holds, then our monomial is proportional to $\widetilde{u}_0 \cdots \widetilde{u}_j \widetilde{I}$ for \widetilde{I} a complete contraction of $\widetilde{\nabla}^{n/2} \widetilde{\operatorname{Rm}}$. Otherwise $S_{i_0} \leq n+1$. In either case, Fefferman and Graham showed [4, Propositions 6.2 and 9.1] that the contribution of the factor $\widetilde{\nabla}^{M_{i_0}} \widetilde{\operatorname{Rm}}$ is independent of the ambiguity of \widetilde{g} . We conclude that our monomial is independent of the ambiguity of \widetilde{g} .

Suppose now that $\ell \geq j+2$ and $k \leq n/2+1$. If $i_0 \in \{1,\ldots,j\}$, then Equation (2.5) yields

$$S_{i_0} \le \sum_{i=1}^{\ell} S_i - 4(\ell - j) \le 2(k - \ell + j) \le 2k - 4 \le n - 2.$$

Lemma 2.2 then implies that the contribution of the factor $\widetilde{\nabla}^{M_{i_0}}\widetilde{u}_{i_0}$ is independent of the ambiguity of \widetilde{g} . If instead $i_0 \in \{j+1,\ldots,\ell\}$, then Equation (2.5) yields

$$S_{i_0} \le \sum_{i=1}^{\ell} S_i - 4(\ell - j - 1) \le 2(k + 2 - \ell + j) \le 2k \le n + 2.$$

Moreover, equality holds if and only if $S_i = 0$ for each $i \leq j$ and $S_i = 4$ for each $i \in \{j+1,\ldots,\ell\}\setminus\{i_0\}$. Fefferman and Graham showed [4, Propositions 6.2 and 9.1] that the contribution of the factor $\widetilde{\nabla}^{M_{i_0}}$ $\widetilde{\operatorname{Rm}}$ is independent of the ambiguity of \widetilde{g} . We again conclude that our monomial is independent of the ambiguity of \widetilde{g} .

Now let $D \colon \mathcal{E}\left[-\frac{n-2k}{j+1}\right]^{\otimes j} \to \mathcal{E}\left[-\frac{jn+2k}{j+1}\right]$ be a conformally invariant polydifferential operator. Then for every compact conformal manifold (M^n,\mathfrak{c}) , the Dirichlet

form $\mathfrak{D} \colon \mathcal{E} \big[- \frac{n-2k}{j+1} \big]^{\otimes (j+1)} \to \mathbb{R}$ determined by

$$\mathfrak{D}(u_0 \otimes \cdots \otimes u_j) := \int_M u_0 D(u_1 \otimes \cdots \otimes u_j) \, dV,$$

is conformally invariant. We say that D is **formally self-adjoint** if \mathfrak{D} is symmetric. This implies that D is itself symmetric.

We conclude this section by constructing the curved Ovsienko–Redou operators D_{2k} and their linear analogues $D_{2k,\mathcal{I}}$. To that end, we identify a tangential linear combination of operators of the form $\widetilde{\Delta}^{k-j} \circ \widetilde{f} \circ \widetilde{\Delta}^j$ for some fixed homogeneous function $\widetilde{f} \in \widetilde{\mathcal{E}}[w']$ (cf. [2, Lemma 6.1]).

Lemma 2.4. Let $(\widetilde{\mathcal{G}}, \widetilde{g})$ be an ambient space for a conformal manifold (M^n, \mathfrak{c}) and let $\widetilde{f} \in \widetilde{\mathcal{E}}[w']$. Let $k \in \mathbb{N}$ and $w \in \mathbb{R}$. Define $\widetilde{D}_{2k,w,\widetilde{f}} \colon \widetilde{\mathcal{E}}[w] \to \widetilde{\mathcal{E}}[w+w'-2k]$ by

$$\widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u}) := \sum_{j=0}^{k} a_j \widetilde{\Delta}^{k-j} \left(\widetilde{f} \widetilde{\Delta}^j \widetilde{u} \right),$$

$$a_j := \binom{k}{j} \frac{\Gamma(j+k-w-w'-\frac{n}{2})\Gamma(\frac{n}{2}+w-j)}{\Gamma(k-w-w'-\frac{n}{2})\Gamma(\frac{n}{2}+w-k)}.$$

Then $\iota^*\widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u})$ depends only on $\iota^*\widetilde{u}$, \widetilde{f} , and \widetilde{g} .

Proof. Consider the defining function $Q:=|T|^2$ for $\iota(\mathcal{G})\subseteq\widetilde{\mathcal{G}}$. It suffices to show that the commutator $[\widetilde{D}_{2k,w,\widetilde{f}},Q]=0$ on $\widetilde{\mathcal{E}}[w-2]$, where Q is regarded as a multiplication operator. Graham et al. observed [7, Equation (1.8)] that

$$[\widetilde{\Delta}^\ell,Q]=2\ell\widetilde{\Delta}^{\ell-1}(2T+n+4-2\ell)$$

on $C^{\infty}(\widetilde{\mathcal{G}})$ for any $\ell \in \mathbb{N}$. Therefore

$$[\widetilde{D}_{2k,w,\widetilde{f}},Q] = \sum_{j=0}^{k-1} b_j \widetilde{\Delta}^{k-j-1} \circ \widetilde{f} \circ \widetilde{\Delta}^j$$

on $\widetilde{\mathcal{E}}[w-2]$, where

$$b_i = 2(j+1)(n+2w-2j-2)a_{j+1} - 2(k-j)(2j+2k-2w-2w'-n)a_j$$

Our definition of
$$a_j$$
 yields $b_j = 0$, and hence $[\widetilde{D}_{2k,w,\widetilde{f}},Q] = 0$ on $\widetilde{\mathcal{E}}[w-2]$.

Note that $\widetilde{D}_{k,w,\widetilde{f}}$ is not a differential operator in the sense of Subsection 2.1 because we do not require that \widetilde{f} is an ambient scalar Riemannian invariant. Applying Lemma 2.4 with a suitably chosen ambient scalar Riemannian invariant does produce a tangential different operator:

Corollary 2.5. Let (M^n, \mathfrak{c}) be a conformal manifold and let $\widetilde{I} \in \widetilde{\mathcal{E}}[-2\ell]$ be an ambient scalar Riemannian invariant. Let $k \in \mathbb{N}$; if n is even, then assume additionally that

(i)
$$k \le n/2$$
, if $\ell = 0$; and

(ii)
$$k + \ell \le n/2 + 1$$
, if $\ell \ge 1$.

Then

$$\widetilde{D}_{2k,\widetilde{I}}(\widetilde{u}) := \sum_{r+s=k} \frac{k!}{r!s!} \frac{(\ell+s-1)!(\ell+r-1)!}{(\ell-1)!^2} \widetilde{\Delta}^r \left(\widetilde{I} \widetilde{\Delta}^s \widetilde{u} \right)$$

defines a tangential differential operator $\widetilde{D}_{2k,\widetilde{I}}$: $\widetilde{\mathcal{E}}\left[-\frac{n-2k-2\ell}{2}\right] \to \widetilde{\mathcal{E}}\left[-\frac{n+2k+2\ell}{2}\right]$. In particular, the differential operator

$$D_{2k,\mathcal{I}} \colon \mathcal{E}\left[-\frac{n-2k-2\ell}{2}\right] \to \mathcal{E}\left[-\frac{n+2k+2\ell}{2}\right],$$
$$D_{2k,\mathcal{I}}(\iota^*\widetilde{u}) := (\widetilde{D}_{2k,\widetilde{I}}\widetilde{u}) \circ \iota,$$

is conformally invariant.

Proof. Corollary 2.3 implies that $\widetilde{D}_{2k,\widetilde{I}}$ is independent of the ambiguity of \widetilde{g} . Since $\widetilde{I} \in \widetilde{\mathcal{E}}[-2\ell]$, applying Lemma 2.4 then implies that

$$\widetilde{D}_{2k,\widetilde{I}} = \widetilde{D}_{2k,w,\widetilde{I}}$$

is tangential, where $w := -\frac{n}{2} + k + \ell$.

The curved Ovsienko–Redou operators arise by looking for tangential linear combinations of the operators $\widetilde{D}_{2k-2s} \widetilde{\Lambda}^s \widetilde{f}$.

Corollary 2.6. Let (M^n, \mathfrak{c}) be a conformal manifold. Let $k \in \mathbb{N}$; if n is even, then assume additionally that $k \leq n/2$. Then

$$\begin{split} \widetilde{D}_{2k}(\widetilde{u}\otimes\widetilde{v}) &:= \sum_{r+s+t=k} a_{r,s,t} \widetilde{\Delta}^r \left((\widetilde{\Delta}^s \widetilde{u}) (\widetilde{\Delta}^t \widetilde{v}) \right), \\ a_{r,s,t} &:= \frac{k!}{r!s!t!} \frac{\Gamma\left(\frac{n+4k}{6} - r\right) \Gamma\left(\frac{n+4k}{6} - s\right) \Gamma\left(\frac{n+4k}{6} - t\right)}{\Gamma\left(\frac{n-2k}{6}\right) \Gamma\left(\frac{n+4k}{6}\right)^2}, \end{split}$$

defines a tangential bidifferential operator $\widetilde{D}_{2k} \colon \widetilde{\mathcal{E}}\left[-\frac{n-2k}{3}\right]^{\otimes 2} \to \widetilde{\mathcal{E}}\left[-\frac{2n+2k}{3}\right]$. In particular, the bidifferential operator

$$D_{2k} \colon \mathcal{E}\left[-\frac{n-2k}{3}\right]^{\otimes 2} \to \mathcal{E}\left[-\frac{2n+2k}{3}\right],$$
$$D_{2k}(\iota^*\widetilde{u} \otimes \iota^*\widetilde{v}) := \widetilde{D}_{2k}(\widetilde{u} \otimes \widetilde{v}) \circ \iota$$

is conformally invariant.

Proof. Corollary 2.3 implies that \widetilde{D}_{2k} is independent of the ambiguity of \widetilde{g} . Since $\widetilde{D}_{2k}(\widetilde{u} \otimes \widetilde{v}) = \widetilde{D}_{2k}(\widetilde{v} \otimes \widetilde{u})$, it thus suffices to show that $\widetilde{D}_{2k}(\widetilde{u} \otimes \widetilde{v}) \circ \iota$ depends only on $\iota^*\widetilde{v}$, \widetilde{u} , and \widetilde{g} in order to conclude that \widetilde{D}_{2k} is tangential. Observe that

$$(2.7) \qquad \widetilde{D}_{2k}(\widetilde{u} \otimes \widetilde{v}) = \sum_{s=0}^{k} {k \choose s} \frac{\Gamma(\frac{n+4k}{6} - s)\Gamma(\frac{n-2k}{6} + s)^{2}}{\Gamma(\frac{n-2k}{6})\Gamma(\frac{n+4k}{6})^{2}} \widetilde{D}_{2k-2s, -\frac{n-2k}{3}, \widetilde{\Delta}^{s} \widetilde{u}}(\widetilde{v}).$$

Applying Lemma 2.4 to each $\widetilde{D}_{2k-2s,-\frac{n-2k}{3},\widetilde{\Delta}^s\widetilde{u}}$ implies that $\widetilde{D}_{2k}(\widetilde{u}\otimes\widetilde{v})\circ\iota$ depends only on $\iota^*\widetilde{v}$, \widetilde{u} , and \widetilde{g} .

3. Poincaré spaces

In this section we give an equivalent description of the operators $\widetilde{D}_{2k,w,\widetilde{f}}$ from Lemma 2.4 in terms of the Poincaré space determined by a straight and normal ambient metric. We restrict our attention to the case $w=-\frac{n+w'-2k}{2}$, where w' is the weight of \widetilde{f} , as it is only in this case that the operator on (M^n,g) induced by $\widetilde{D}_{2k,w,\widetilde{f}}$ can be formally self-adjoint.

Let $(\widetilde{\mathcal{G}}, \widetilde{g})$ be the straight and normal ambient space (2.1) for a representative $g \in \mathfrak{c}$ of a conformal manifold (M^n, \mathfrak{c}) . Set $r := \sqrt{-2\rho}$ and s := rt in the domain $\widetilde{\mathcal{G}}_+ := \{\rho < 0, t > 0\} \subseteq \widetilde{\mathcal{G}}$. In these coordinates,

(3.1)
$$\widetilde{g} = -ds^2 + s^2 g_+,$$

$$g_+ = r^{-2} (dr^2 + g_{-r^2/2}).$$

The equation $\widetilde{g}(T,T)=-1$ defines a hypersurface $\mathring{X}\subseteq\widetilde{\mathcal{G}}$. Since $T=t\partial_t$, we see that $\mathring{X}=\{s=1\}$. Using Equation (3.1) to compute the Laplacian $\widetilde{\Delta}$ of \widetilde{g} in terms of the Laplacian of g_+ yields

(3.2)
$$\widetilde{\Delta} = s^{-2} \left(\Delta_{g_+} - (s\partial_s)^2 - ns\partial_s \right).$$

Suppose that $\widetilde{u} \in \widetilde{\mathcal{E}}[w]$. Thus there is a one-parameter family of functions $u_{\rho} \in C^{\infty}(M)$ such that $\widetilde{u}(t,x,\rho) = t^{w}u_{\rho}(x)$. Equivalently, $\widetilde{u}(r,x,s) = s^{w}\widehat{u}(r,x)$, where $\widehat{u} \in C^{\infty}(\mathring{X})$ is the function $\widehat{u}(r,x) = r^{-w}u_{-r^{2}/2}(x)$. In particular, \widetilde{u} is determined by its restriction $\widehat{u} = \widetilde{u}|_{\mathring{X}}$. Moreover, Equation (3.2) implies that

(3.3)
$$\widetilde{\Delta}\widetilde{u} = s^{w-2} \left(\Delta_{q_+} - w(n+w) \right) \widehat{u}.$$

This leads to a formula relating the operators $\widetilde{D}_{2k,w,\widetilde{f}}$ and

$$P_{2j,w} := \prod_{i=0}^{j-1} \left(\Delta_{g_+} - (w-2i)(n+w-2i) \right)$$

on (\mathring{X}, g_+) :

Lemma 3.1. Let $(\widetilde{\mathcal{G}}, \widetilde{g})$ be the straight and normal ambient space for a representative $g \in \mathfrak{c}$ of a conformal manifold (M^n, \mathfrak{c}) . Let (\mathring{X}, g_+) be the Poincaré manifold determined by the variables r, s as in Equation (3.1). Let $\widetilde{f} \in \widetilde{\mathcal{E}}[w']$ and let $k \in \mathbb{N}$. Set $w := -\frac{n+w'-2k}{2}$. If $\widetilde{u} \in \widetilde{\mathcal{E}}[w]$, then

$$\widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u})|_{\widetilde{X}} = \sum_{i=0}^{k} \sum_{j=0}^{k-i} c_{i,j} (P_{2i,w} \circ \widehat{f} \circ P_{2j,w})(\widehat{u}),$$

$$c_{i,j} := 4^{k-i-j} \frac{k!}{i!j!} \frac{\Gamma(\frac{n}{2} + w - i)\Gamma(\frac{n}{2} + w - j)}{\Gamma(\frac{n}{2} + w - k)^2},$$

where $\widehat{u}:=\widetilde{u}|_{\mathring{X}}$ and $\widehat{f}:=\widetilde{f}|_{\mathring{X}}.$

Proof. Note that w + w' - k = k - n - w. Lemma 2.4 and Equation (3.3) yield

$$(3.4) \quad \widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u})|_{\widetilde{X}} = \sum_{j=0}^{k} {k \choose j} \frac{\Gamma(\frac{n}{2} + w - k + j)\Gamma(\frac{n}{2} + w - j)}{\Gamma(\frac{n}{2} + w - k)^{2}} \times (P_{2k-2j,2k-2j-n-w} \circ \widehat{f} \circ P_{2j,w})(\widehat{u}).$$

On the one hand, a simple reindexing in the definition of $P_{2j,w}$ yields

$$(3.5) P_{2k-2j,2k-2j-n-w} = P_{2k-2j,w-2}.$$

On the other hand, direct computation yields

$$\Delta_{g_{+}} - (w - 2j)(n + w - 2j) = \Delta_{g_{+}} - w(n + w) + 2j(n + 2w - 2j).$$

It follows that

$$P_{2j,w-2} = P_{2j,w} + 4j\left(\frac{n}{2} + w - j\right)P_{2j-2,w-2}.$$

A simple induction argument yields

(3.6)
$$P_{2j,w-2} = \sum_{i=0}^{j} 4^{i} \frac{j!}{(j-i)!} \frac{\Gamma(\frac{n}{2} + w - j + i)}{\Gamma(\frac{n}{2} + w - j)} P_{2j-2i,w}.$$

Combining Equations (3.4), (3.5), and (3.6) yields

$$\widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u})|_{X}^{*} = \sum_{j=0}^{k} \sum_{i=0}^{k-j} 4^{i} \frac{k!}{j!(k-i-j)!} \frac{\Gamma(\frac{n}{2} + w - k + i + j)\Gamma(\frac{n}{2} + w - j)}{\Gamma(\frac{n}{2} + w - k)^{2}} \times (P_{2k-2i-2j,w} \circ \widehat{f} \circ P_{2j,w})(\widehat{u}).$$

Reindexing the sum yields the final result.

4. Formal self-adjointness

We are now ready to prove the formal self-adjointness of the curved Ovsienko–Redou operators D_{2k} and the conformally invariant operators $D_{2k,\mathcal{I}}$. To that end, we identify their associated Dirichlet forms as the logarithmic term in the expansion of the corresponding Dirichlet form in a Poincaré space. More generally:

Lemma 4.1. Let $(\widetilde{\mathcal{G}}, \widetilde{g})$ be the straight and normal ambient space for a representative $g \in \mathfrak{c}$ of a conformal manifold (M^n, \mathfrak{c}) . Denote by (\mathring{X}, g_+) the Poincaré space determined by the variables r, s as in Equation (3.1). Let $\widetilde{f} \in \widetilde{\mathcal{E}}[w']$ and let $k \in \mathbb{N}$. Set $w := -\frac{n+w'-2k}{2}$. Define $D_{2k,w,\widetilde{f}} \colon C^{\infty}(M) \to C^{\infty}(M)$ by

$$D_{2k,w,\tilde{f}}(u) := \tilde{D}_{2k,w,\tilde{f}}(t^w \pi^* u)|_{t=1,\rho=0}.$$

If $\widetilde{u}, \widetilde{v} \in \widetilde{\mathcal{E}}[w]$, then

$$\int_{r>\varepsilon} \left. \left(\widetilde{v} \widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u}) \right) \right|_{\mathring{X}} \mathrm{dV}_{g_+} = - \left(\int_{M} v D_{2k,w,\widetilde{f}}(u) \, \mathrm{dV}_g \right) \log \varepsilon + O(1)$$

as $\varepsilon \to 0$, where $u := \widetilde{u}|_{t=1,\rho=0}$ and $v := \widetilde{v}|_{t=1,\rho=0}$.

Proof. Note that 2w + w' - 2k = -n. Thus $\widetilde{v}\widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u}) \in \widetilde{\mathcal{E}}[-n]$. Therefore

$$\left(\widetilde{v}\widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u})\right)|_{\mathring{X}} = vD_{2k,w,\widetilde{f}}(u)r^n + O(r^{n+2})$$

as $r \to 0$. Graham observed [6, Equations (3.1) and (3.2)] that

$$dV_{q_{+}} = r^{-n-1} (1 + O(r^{2})) dV_{q} dr$$

as $r \to 0$. Direct calculation yields the final result.

Combining Lemmas 3.1 and 4.1 implies that $D_{2k,w,\tilde{t}}$ is formally self-adjoint.

Proposition 4.2. Let $(\widetilde{\mathcal{G}}, \widetilde{g})$ be the straight and normal ambient space for a representative $g \in \mathfrak{c}$ of a conformal manifold (M^n, \mathfrak{c}) . Denote by (\mathring{X}, g_+) the Poincaré space as in Equation (3.1). Let $\widetilde{f} \in \widetilde{\mathcal{E}}[w']$ and let $k \in \mathbb{N}$. Set $w := -\frac{n+w'-2k}{2}$. Define $D_{2k,w,\widetilde{f}} \colon C^{\infty}(M) \to C^{\infty}(M)$ as in Lemma 4.1. Then

$$\int_{M} v D_{2k,w,\widetilde{f}} u \, \mathrm{dV}_g = \int_{M} u D_{2k,w,\widetilde{f}}(v) \, \mathrm{dV}_g$$

for every compactly supported $u, v \in C^{\infty}(M)$.

Proof. Let $u, v \in C^{\infty}(M)$ be compactly supported. Set $\widetilde{u} := t^w \pi^* u$ and $\widetilde{v} := t^w \pi^* v$. Given a function $\phi = \phi(\varepsilon)$ such that $\phi \in O(\log \varepsilon)$ as $\varepsilon \to 0$, denote

$$\operatorname{lp} \phi := \lim_{\varepsilon \to 0} \frac{\phi(\varepsilon)}{\log(1/\varepsilon)}.$$

First, Lemma 4.1 yields

$$(4.1) \int_{M} \left(v D_{2k,w,\widetilde{f}}(u) - u D_{2k,w,\widetilde{f}}(v) \right) dV_{g}$$

$$= \ln \int_{T \setminus \widetilde{s}} \left(\widetilde{v} \widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u}) - \widetilde{u} \widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{v}) \right) \Big|_{\mathring{X}} dV_{g_{+}}.$$

Second, Lemma 3.1 and the symmetry $c_{i,j} = c_{j,i}$ yield

$$(4.2) \int_{r>\varepsilon} \left(\widetilde{v} \widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{u}) - \widetilde{u} \widetilde{D}_{2k,w,\widetilde{f}}(\widetilde{v}) \right) \Big|_{\mathring{X}} dV_{g_{+}}$$

$$= \sum_{i+j=0}^{k} c_{i,j} \int_{r>\varepsilon} \left(\widehat{v} (P_{2i,w} \circ \widehat{f} \circ P_{2j,w}) (\widehat{u}) - \widehat{u} (P_{2j,w} \circ \widehat{f} \circ P_{2i,w}) (\widehat{v}) \right) dV_{g_{+}},$$

where $\widehat{u} := \widetilde{u}|_{\mathring{X}}$ and $\widehat{v} := \widetilde{v}|_{\mathring{X}}$. Third, observe that if $0 \leq j \leq i$, then

$$P_{2i,w} = P_{2i-2j,w-2j} \circ P_{2j,w}$$
.

Let $\widehat{\varphi}$, $\widehat{\psi} \in C^{\infty}(\mathring{X})$. Write

$$\begin{split} & \int_{r>\varepsilon} \left(\widehat{\varphi} P_{2i,w} \widehat{\psi} - \widehat{\psi} P_{2i,w} \widehat{\varphi} \right) \mathrm{dV}_{g_+} \\ & = \sum_{\ell=1}^i \int_{r>\varepsilon} \left((P_{2i-2\ell,w-2\ell} \widehat{\varphi}) P_{2\ell,w} \widehat{\psi} - (P_{2i-2\ell+2,w-2\ell+2} \widehat{\varphi}) P_{2\ell-2,w} \widehat{\psi} \right) \mathrm{dV}_{g_+} \\ & = \sum_{\ell=1}^i \int_{r>\varepsilon} \left((P_{2i-2\ell,w-2\ell} \widehat{\varphi}) \Delta_{g_+} P_{2\ell-2,w} \widehat{\psi} - (\Delta_{g_+} P_{2i-2\ell,w-2\ell} \widehat{\varphi}) P_{2\ell-2,w} \widehat{\psi} \right) \mathrm{dV}_{g_+} \,. \end{split}$$

The Divergence Theorem yields

$$\begin{split} & \int_{r>\varepsilon} \left(\widehat{\varphi} P_{2i,w} \widehat{\psi} - \widehat{\psi} P_{2i,w} \widehat{\varphi} \right) \mathrm{dV}_{g_{+}} \\ & = \sum_{\ell=1}^{i} \int_{r=\varepsilon} \left((P_{2i-2\ell,w-2\ell} \widehat{\varphi}) \nabla_{\mathsf{n}} P_{2\ell-2,w} \widehat{\psi} - (\nabla_{\mathsf{n}} P_{2i-2\ell,w-2\ell} \widehat{\varphi}) P_{2\ell-2,w} \widehat{\psi} \right) \mathrm{dA}_{g_{+}}, \end{split}$$

where \mathbf{n} and dA_{g_+} are the outward-pointing unit normal and the area element, respectively, of $\{r = \varepsilon\}$ with respect to g_+ . In particular,

$$\operatorname{lp} \int_{r>\varepsilon} \left(\widehat{\varphi} P_{2i,w} \widehat{\psi} - \widehat{\psi} P_{2i,w} \widehat{\varphi} \right) dV_{g_{+}} = 0.$$

Combining this with Equation (4.1) and (4.2) yields the final result.

Our main results follow from Proposition 4.2 and tangentiality.

Proof of Theorem 1.1. By Corollary 2.6, it suffices to compute in the straight and normal ambient space associated to a representative $g \in \mathfrak{c}$. Equation (2.7) and Proposition 4.2 express D_{2k} as a linear combination of formally self-adjoint operators. Therefore D_{2k} is formally self-adjoint.

Proof of Theorem 1.2. By Corollary 2.5, it suffices to compute in the straight and normal ambient space associated to a representative $g \in \mathfrak{c}$. Combining Equation (2.6) with Proposition 4.2 realizes $D_{2k,\mathcal{I}}$ as a formally self-adjoint operator. \square

Acknowledgements

We would like to thank Shane Chern for very useful discussions.

JSC acknowledges support from a Simons Foundation Collaboration Grant for Mathematicians, ID 524601.

REFERENCES

- T. N. Bailey, M. G. Eastwood, and C. R. Graham, Invariant theory for conformal and CR geometry, Ann. of Math. (2) 139 (1994), no. 3, 491–552. MR1283869
- [2] J. S Case, Y.-J. Lin, and W. Yuan, Curved Versions of the Ovsienko-Redou Operators, Int. Math. Res. Not. IMRN 19 (2023), 16904–16929. MR4651902
- [3] C. Fefferman and C. R. Graham, Q-curvature and Poincaré metrics, Math. Res. Lett. 9 (2002), no. 2-3, 139–151. MR1909634
- [4] ______, The ambient metric, Annals of Mathematics Studies, vol. 178, Princeton University Press, Princeton, NJ, 2012. MR2858236
- [5] _____, Juhl's formulae for GJMS operators and Q-curvatures, J. Amer. Math. Soc. 26 (2013), no. 4, 1191–1207. MR3073887
- [6] C. R. Graham, Volume and area renormalizations for conformally compact Einstein metrics, The Proceedings of the 19th Winter School "Geometry and Physics" (Srní, 1999), 2000, pp. 31–42. MR1758076
- [7] C. R. Graham, R. Jenne, L. J. Mason, and G. A. J. Sparling, Conformally invariant powers of the Laplacian. I. Existence, J. London Math. Soc. (2) 46 (1992), no. 3, 557–565. MR1190438
- [8] C. R. Graham and M. Zworski, Scattering matrix in conformal geometry, Invent. Math. 152 (2003), no. 1, 89–118. MR1965361
- [9] A. Juhl, Explicit formulas for GJMS-operators and Q-curvatures, Geom. Funct. Anal. 23 (2013), no. 4, 1278–1370. MR3077914
- [10] V. Ovsienko and P. Redou, Generalized transvectants-Rankin-Cohen brackets, Lett. Math. Phys. 63 (2003), no. 1, 19–28. MR1967533

Department of Mathematics, Penn State University, University Park, PA 16802, USA $\it Email~address:$ <code>jscase@psu.edu</code>

DEPARTMENT OF MATHEMATICS, UC SANTA BARBARA, SANTA BARBARA, CA 93106, USA

Email address: ztyan@ucsb.edu